

**Department of Radiology**  
Interventional Neuro-Radiology Discharge  
Instructions for Aneurysm Embolization

1. May return to non-strenuous work in 2-4 days after discharge with the following restrictions:
  - Weight limit of 10 lbs.
  - No strenuous activity until seen in clinic.
  - Additional instructions: \_\_\_\_\_
  
2. Resume previously prescribed medication and begin new medications as directed.
3. Take Tylenol as directed for discomfort or headache.
4. **Do not take Aspirin, Motrin, Naprosyn or any medication containing these ingredients for 6 weeks unless instructed otherwise.**
5. You will be notified to schedule your next procedure or clinic visit.

**NOTIFY DR LARSON'S OFFICE AT 615-322-3357, OR AFTER HOURS AND WEEKENDS AT 615 322-5500 (Ask the operator to page Dr. Larson) FOR ANY OF THE FOLLOWING SYMPTOMS:**

- Fever greater than 101° F orally.
- Any stroke-like symptoms, i.e. Changes in mental status, memory or vision, weakness, numbness, tingling, or pain in arms, legs or face, or difficulty speaking or swallowing.
- Severe or unusual headache
- Seizure

**HOME CARE INSTRUCTIONS FOR PUNCTURE SITE**

- If bleeding occurs from puncture site, apply direct pressure for 15 minutes.
  - If bleeding stops after 15 minutes, lie flat for 2 hours.
  - If bleeding continues, hold pressure and either call 911 or have someone drive you to the nearest Emergency Room.
- Notify us if you have more than minor bleeding, pain, redness or discharge at the puncture site.
- Bathing of puncture site is permitted after 48 hours but dry the site well.

\_\_\_\_\_  
Patient or Responsible Adult

\_\_\_\_\_  
Date/Time

\_\_\_\_\_  
Nurse

\_\_\_\_\_  
Date/Time



**Department of Radiology**  
Interventional Neuro-Radiology Discharge  
Instructions for Aneurysm Embolization

1. May return to non-strenuous work in 2-4 days after discharge with the following restrictions:
  - Weight limit of 10 lbs.
  - No strenuous activity until seen in clinic.
  - Additional instructions: \_\_\_\_\_
  
2. Resume previously prescribed medication and begin new medications as directed.
3. Take Tylenol as directed for discomfort or headache.
4. **Do not take Aspirin, Motrin, Naprosyn or any medication containing these ingredients for 6 weeks unless instructed otherwise.**
5. You will be notified to schedule your next procedure or clinic visit.

**NOTIFY DR LARSON'S OFFICE AT 615-322-3357, OR AFTER HOURS AND WEEKENDS AT 615 322-5500 (Ask the operator to page Dr. Larson) FOR ANY OF THE FOLLOWING SYMPTOMS:**

- Fever greater than 101° F orally.
- Any stroke-like symptoms, i.e. Changes in mental status, memory or vision, weakness, numbness, tingling, or pain in arms, legs or face, or difficulty speaking or swallowing.
- Severe or unusual headache
- Seizure

**HOME CARE INSTRUCTIONS FOR PUNCTURE SITE**

- If bleeding occurs from puncture site, apply direct pressure for 15 minutes.
  - If bleeding stops after 15 minutes, lie flat for 2 hours.
  - If bleeding continues, hold pressure and either call 911 or have someone drive you to the nearest Emergency Room.
- Notify us if you have more than minor bleeding, pain, redness or discharge at the puncture site.
- Bathing of puncture site is permitted after 48 hours but dry the site well.

\_\_\_\_\_  
Patient or Responsible Adult

\_\_\_\_\_  
Date/Time

\_\_\_\_\_  
Nurse

\_\_\_\_\_  
Date/Time